

Dynamical Electroweak Symmetry Breaking in $SO(5) \times U(1)$ Gauge-Higgs Unification in the Randall-Sundrum Warped Space

Yutaka Hosotani

Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan

Abstract

In the gauge-Higgs unification scenario the Higgs field is unified with gauge fields in higher dimensional gauge theory. The 4D Higgs field $H(x)$ corresponds to 4D fluctuations of the Aharonov-Bohm phase (Wilson line phase) θ_H in the extra-dimension. An $SO(5) \times U(1)$ gauge-Higgs unification model in the Randall-Sundrum warped spacetime with top and bottom quarks is presented. Gauge couplings of the top quark multiplet induce electroweak symmetry breaking by the Hosotani mechanism. The effective potential $V_{\text{eff}}(\theta_H)$ is found to be minimized at $\theta_H = \frac{1}{2}\pi$ and the Higgs mass is predicted around 50 GeV. The ZZH and WWH couplings vanish at $\theta_H = \frac{1}{2}\pi$ so that the LEP2 bound for the Higgs mass is evaded. The result is summarized in the effective interactions for $\hat{\theta}_H(x) = \theta_H + H(x)/f_H$.

Key words: gauge-Higgs unification, electroweak symmetry breaking, Hosotani mechanism

PACS: 11.10.Kk, 11.15.Ex, 12.60.-i, 12.60.Cn

OU-HET 621/2009

1. Introduction

In the standard model the electroweak (EW) symmetry is spontaneously broken by the Higgs mechanism, which is yet to be confirmed by experiments. Although the Higgs particle is expected to be found at LHC in the near future, it is not clear whether or not the Higgs particle appears as described in the standard model.

Recently there has been significant progress in the gauge-Higgs unification scenario in which the 4D Higgs field is identified with a part of the extra-dimensional component of gauge fields in higher dimensions. The Higgs field appears as an Aharonov-Bohm (AB) phase, or a Wilson line phase, in the extra dimension, thereby the EW symmetry being dynamically broken by the Hosotani mechanism.[1,2] One of the promising models is




Fig. 1. The effective potential $V_{\text{eff}}(\theta_H)$. $U(\theta_H/\pi) = (4\pi)^2(kz_L^{-1})^{-4}V_{\text{eff}}(\theta_H)$ is plotted for $z_L = 10^{15}$.

the $SO(5) \times U(1)_X$ gauge-Higgs unification model in the Randall-Sundrum (RS) warped spacetime, in which many predictions are given.[3]-[9]

2. Model

The metric of the RS warped spacetime is given by $ds^2 = z^{-2}(\eta_{\mu\nu}dx^\mu dx^\nu + k^{-2}dz^2)$, $1 \leq z \leq z_L = e^{kL}$. The bulk region $1 < z < z_L$ is the AdS space with a cosmological constant $\Lambda = -6k^2$, which is sandwiched by the Planck brane at $z = 1$ and the TeV brane at $z = z_L$. We consider an $SO(5) \times U(1)_X$ gauge theory in this space.[7] The orbifold boundary conditions at the two branes break the symmetry to $SO(4) \times U(1)_X$. Additional brane interactions on the Planck brane further break it down to the EW symmetry $SU(2)_L \times U(1)_Y$. The 4D $SU(2)_L$ doublet Higgs field resides in the $(a5)$ -components ($a = 1, \dots, 4$) of $A_z(x, z)$. The Aharonov-Bohm (AB) phase is given by the phase of $(P \exp \{ig \int_1^{z_L} dz A_z\})^2$. $A_z^{(45)}$ contains the 4D neutral Higgs field $H(x)$.

Fermions are introduced in the vector $(\mathbf{5})$ representation of $SO(5)$ in the bulk. In terms of $SU(2)_L$ content they are, for the top and bottom multiplets,

$$\left(\begin{bmatrix} T \\ B \end{bmatrix}, \begin{bmatrix} t \\ b \end{bmatrix}, t' \right), \quad \left(\begin{bmatrix} U \\ D \end{bmatrix}, \begin{bmatrix} X \\ Y \end{bmatrix}, b' \right). \quad (1)$$

In addition, right-handed brane fermions in the $(\mathbf{2}, \mathbf{1})$ representation of $SU(2)_L \times SU(2)_R$ are introduced on the Planck brane. At low energies the desired mass spectrum for the top and bottom quarks is obtained. All other fermions become heavy.

When the AB phase θ_H associated with $A_z^{(45)}$ becomes non-zero, the EW symmetry is broken, thereby W , Z , top and bottom quarks acquiring masses. In particular,

$$m_t \sim \frac{m_{\text{KK}}}{\sqrt{2}\pi} \sqrt{1 - 4c^2} |\sin \theta_H|, \quad m_b \sim \left| \frac{\tilde{\mu}}{\mu_2} \right| m_t \quad (2)$$

where the Kaluza-Klein mass scale is given by $m_{\text{KK}} \sim \pi k z_L^{-1} \sim 1.5 \text{ TeV}$. c is the bulk mass parameter (~ 0.43). $\tilde{\mu}$ and μ_2 are mass couplings between the bulk and brane fermions.

3. Dynamical EW symmetry breaking

The effective potential $V_{\text{eff}}(\theta_H)$ is evaluated at the one loop level.[6,7,10,11,12] The result is depicted in fig. 1. Without the fermions $V_{\text{eff}}(\theta_H)$ is minimized at $\theta_H = 0$ or π so that the EW gauge symmetry is unbroken. With the contributions from the top quark multiplet, V_{eff} has global minima at $\theta_H = \pm \frac{1}{2}\pi$ so that the EW gauge symmetry is dynamically broken. Contributions from light quarks (c, s, u, d) are negligible. In the flat spacetime limit ($k \rightarrow 0$) the symmetry remains unbroken.

The Higgs mass is given by

$$m_H^2 = \frac{1}{f_H^2} \left. \frac{d^2 V_{\text{eff}}}{d\theta_H^2} \right|_{\min}, \quad f_H \sim \frac{2}{\sqrt{kL}} \frac{m_{\text{KK}}}{\pi g}. \quad (3)$$

m_H is predicted around 50 GeV for $z_L = 10^{13} \sim 10^{17}$. We stress that this is in no conflict with the LEP2 bound for m_H . The ZZH coupling vanishes at $\theta_H = \frac{1}{2}\pi$. The process $e^+e^- \rightarrow Z \rightarrow ZH$ does not take place so that the LEP2 bound is evaded.

4. Effective interactions

The Higgs interactions are summarized in the effective interactions[5,9]

$$\begin{aligned} \mathcal{L}_{\text{eff}} &= -V_{\text{eff}}(\hat{\theta}_H) - m_W(\hat{\theta}_H)^2 W_\mu^\dagger W^\mu - \frac{1}{2} m_Z(\hat{\theta}_H)^2 Z_\mu Z^\mu - \sum_f m_f(\hat{\theta}_H) \bar{\psi}_f \psi_f, \\ m_W(\hat{\theta}_H) &\sim \cos \theta_W m_Z(\hat{\theta}_H) \sim \frac{1}{2} g f_H \sin \hat{\theta}_H, \\ m_f(\hat{\theta}_H) &\sim \lambda_f \sin \hat{\theta}_H, \quad \hat{\theta}_H(x) = \theta_H + \frac{H(x)}{f_H}. \end{aligned} \quad (4)$$

The first term is the effective potential for $\hat{\theta}_H$. $m_W(\hat{\theta}_H)$ and $m_Z(\hat{\theta}_H)$ are given in [4]. $m_W = m_W(\theta_H)$, $m_Z = m_Z(\theta_H)$. Expanding $m_W(\hat{\theta}_H)^2$ and $m_Z(\hat{\theta}_H)^2$ in a power series in H , one finds that the WWH and ZZH couplings are suppressed by a factor $\cos \theta_H$ compared with those in the standard model. For the $WWHH$ and $ZZHH$ couplings the suppression factor becomes $\cos 2\theta_H$. $m_f(\hat{\theta}_H)$ is given in [9]. It follows that the WWH , ZZH and Yukawa couplings vanish at $\theta_H = \frac{1}{2}\pi$, which leads to new phenomenology.

References

- [1] Y. Hosotani, *Phys. Lett.* **B126** (1983) 309; *Ann. Phys. (N.Y.)* **190** (1989) 233.
- [2] A.T. Davies and A. McLachlan, *Phys. Lett.* **B200** (1988) 305; *Nucl. Phys.* **B317** (1989) 237.
- [3] K. Agashe, R. Contino and A. Pomarol, *Nucl. Phys.* **B719** (2005) 165.
- [4] Y. Sakamura and Y. Hosotani, *Phys. Lett.* **B645** (2007) 442; *Prog. Theoret. Phys.* **118** (2007) 935.
- [5] Y. Sakamura, *Phys. Rev.* **D76** (2007) 065002.
- [6] A.D. Medina, N.R. Shah and C.E.M. Wagner, *Phys. Rev.* **D76** (2007) 095010.
- [7] Y. Hosotani, K. Oda, T. Ohnuma and Y. Sakamura, *Phys. Rev.* **D78** (2008) 096002.
- [8] Y. Hosotani, arXiv:0809.2181[hep-ph].
- [9] Y. Hosotani and Y. Kobayashi, arXiv:0812.4782[hep-ph].
- [10] K. Oda and A. Weiler, *Phys. Lett.* **B606** (2005) 408.

- [11] A. Falkowski, *Phys. Rev. D* **75** (2007) 025017.
- [12] H. Hatanaka, arXiv: 0712.1334 [hep-th].